

## True (T) or False (F):

### Regarding South African adolescents' knowledge of abortion legislation, and attitude to abortion

1. The majority of adolescents considered abortion an acceptable option for pregnancy.
2. Only 10% of adolescents were aware of the period of gestation when termination is allowed.
3. Information on the basic aspects of reproductive rights should be central to Life Orientation classes in schools.

### Regarding adolescent preventive services

4. Anticipatory guidance and preventive services should be offered to adolescents during routine clinical practice.
5. Adolescents make up one-fifth of the population in Nigeria.

### Regarding iron deficiency in primary school learners

6. Iron deficiency anaemia is associated with a high serum ferritin level.
7. In primary school learners in the Pretoria region, iron deficiency is the most common cause of anaemia.
8. Children with iron deficiency anaemia perform less well than controls in learning areas such as life skills and language.

### Regarding the early diagnosis and treatment (EIDT) of HIV in infants

9. In 2013 it was estimated that ~25% of children <15 years old living with HIV in sub-Saharan Africa were on antiretroviral therapy.
10. EIDT aims to identify and test all HIV-exposed infants and commence treatment of those infected by 24 months of age.

### Regarding dermatological conditions seen in a specialised outpatient clinic in KwaZulu-Natal

11. Two-thirds of children referred to a dermatology clinic have the correct diagnosis made by the referring doctor.
12. Infective skin conditions are the most common cause for referral.

### Regarding cochlear implantation in South Africa

13. Globally ~1:175 babies are born with hearing impairment.
14. Cochlear implantation is only available in the private health sector in SA.

### Regarding dysphagia in neonates

15. Tube feeding is an effective management of oropharyngeal dysphagia in neonates.
16. A modified barium swallow is considered an appropriate diagnostic aid in the diagnosis of dysphagia in neonates.

### Regarding short-stay ward admissions in the Cape Peninsula

17. Preventable or treatable diseases make up 75% of all under-5 deaths in South Africa.
18. The Western Cape has the least variation in under-5 mortality between regions of all the provinces in South Africa.

### Regarding infant hearing screening

19. Formalised hearing tests in infants are conducted in the majority of primary healthcare clinics in Gauteng.
20. In planning early hearing detection and intervention programmes, district health structures need to be consulted and involved.

A maximum of 3 CEUs will be awarded per correctly completed test.

CPD questionnaires must be completed online via [www.mpconsulting.co.za](http://www.mpconsulting.co.za). After submission you can check the answers and print your certificate. Accreditation number: MDB015/177/02/2016 (Clinical)

